



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GEOMETRY.

151. Proposed by FRANK A. GRIFFIN, Assistant in Mathematics, University of Colorado, Boulder, Col.

A point is at a distance of 1 inch, 2 inches, and $2\frac{1}{2}$ inches, respectively, from three corners of a square. Construct the square. Also solve for the general distances a , b , c .

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa.; P. S. BERG, B. Sc., Larimore, N. D.; G. W. DROKE, State University, Lafayette, Ark.; and MARTIN SPINKS, Wilmington, Ohio.

Let P be the point, $PA=a$, $PB=b$, $PC=c$, $AB=x$ =side of square, $PE=m$, $PF=n$, $m^2+n^2=b^2$(1).

$$a^2-(x+m)^2=a^2-x^2-2mx-m^2=n^2\ldots(2).$$

$$c^2-(x+n)^2=c^2-x^2-2nx-n^2=m^2\ldots(3).$$

Adding (2) and (3) and using (1) we get

$$n+m=\frac{a^2+c^2-2b^2-2x^2}{2x}\ldots(4).$$

Subtracting (3) from (2) we get

$$n-m=\frac{c^2-a^2}{2x}\ldots(5). \quad \therefore n=\frac{c^2-b^2-x^2}{2x}, \quad m=\frac{a^2-b^2-x^2}{2x}.$$

These values of m and n in (1) give

$$x^4-(a^2+c^2)x^2=b^2(a^2+c^2)-\frac{a^4+c^4+2b^4}{2}.$$

$$\therefore x=\left\{\frac{1}{2}[a^2+c^2\pm\sqrt{4b^2(a^2+c^2-b^2)-(a^2-c^2)^2}]\right\}^{\frac{1}{2}}$$

=1.5163 inches when P is without the square.
=2.8197 inches when P is within the square.

I. When P is without, lay off $PE=m$, perpendicular to PE lay off $EB=n$ and $BC=x$, also AB perpendicular to $EB=x$. The square is now determined.

II. Similarly, when P is within, only $EC=x-n$.

152. Proposed by ELMER SCHUYLER, Reading, Pa.

Find a point in a given straight line such that tangents drawn from it to two given circles shall make equal angles with the line.

Solution by MARCUS BAKER, Washington, D. C.

In the annexed figure let A be the given line, and C and c the given circles.

Regarding line A as an axis, revolve C and c about it until they fall at C' and c' .

Then draw the four common tangents 1, 2, 3, and 4 of circles C and c' intersecting the axis in I, II, III, and IV. Each of these points fulfils the conditions. The proof is obvious.

Also solved by G. B. M. ZERR, and D. B. NORTHROP.

